

# UNCLASSIFIED

<b>AD NUMBER</b>
ADB248567
<b>NEW LIMITATION CHANGE</b>
<b>TO</b> Approved for public release, distribution unlimited
<b>FROM</b> Distribution authorized to U.S. Gov't. agencies only; Proprietary Info; Jun 99. Other requests shall be referred to U.S. Army Medical Research and Materiel Command, 504 Scott Street, Ft. Detrick, MD 21702-5012.
<b>AUTHORITY</b>
US Army Med Research and Mat Cmd, MCMR-RMI-S [70-1y], ltr 6 Jul 2000, Ft Detrick, MD

THIS PAGE IS UNCLASSIFIED

AD \_\_\_\_\_

Grant Number DAMD17-96-1-6073

TITLE: The Identification of Estrogen Responsive Genes in the Hypothalamic-Pituitary Axis

PRINCIPAL INVESTIGATOR: Caroline E. Connor

CONTRACTING ORGANIZATION: Duke University Medical Center  
Durham, North Carolina 27710

REPORT DATE: June 1999

TYPE OF REPORT: Annual Summary

PREPARED FOR: U.S. Army Medical Research and Materiel Command  
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Distribution authorized to U.S. Government agencies only (proprietary information, Jun 99). Other requests for this document shall be referred to U.S. Army Medical Research and Materiel Command, 504 Scott Street, Fort Detrick, Maryland 21702-5012.

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

19991027 049

# REPORT DOCUMENTATION PAGE

Form Approved

OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

<b>1. AGENCY USE ONLY (Leave blank)</b>		<b>2. REPORT DATE</b> June 1999		<b>3. REPORT TYPE AND DATES COVERED</b> Annual Summary (1 Jul 96 - 30 Jun 99)	
<b>4. TITLE AND SUBTITLE</b> The Identification of Estrogen Responsive Genes in the Hypothalamic-Pituitary Axis				<b>5. FUNDING NUMBERS</b> DAMD17-96-1-6073	
<b>6. AUTHOR(S)</b> Caroline E. Connor					
<b>7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)</b> Duke University Medical Center Durham, North Carolina 27710				<b>8. PERFORMING ORGANIZATION REPORT NUMBER</b>	
<b>9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)</b> U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012				<b>10. SPONSORING/MONITORING AGENCY REPORT NUMBER</b>	
<b>11. SUPPLEMENTARY NOTES</b>					
<b>12a. DISTRIBUTION / AVAILABILITY STATEMENT</b> Distribution authorized to U.S. Government agencies only (proprietary information, Jun 99). Other requests for this document shall be referred to U.S. Army Medical Research and Materiel Command, 504 Scott Street, Fort Detrick, Maryland 21702-5012.				<b>12b. DISTRIBUTION CODE</b>	
<b>13. ABSTRACT (Maximum 200)</b> <p>Tamoxifen, the most widely used adjuvant therapy for the treatment of estrogen receptor-positive breast cancer, has been shown to inhibit the growth of MCF-7 breast cancer cells in athymic mice. However, in both the animal model and clinical setting, therapeutic utility is limited by the development of resistance to the drug. Our goal is to determine whether GW5638, a structurally similar yet mechanistically distinct compound, can inhibit the growth of tamoxifen-refractory tumors.</p> <p>We demonstrated that GW5638 was able to inhibit the growth of estrogen-dependent MCF-7 tumors in athymic mice. We subsequently developed tumor variants which are no longer sensitive to the inhibitory effects of tamoxifen; the tumors in fact require tamoxifen, but not estrogen, for growth. The resistant tumors were implanted into athymic mice treated with tamoxifen, GW5638, or both drugs. Two important findings emerged from these studies: 1) GW5638 alone cannot support the growth of tamoxifen-dependent tumors, and 2) GW5638 inhibits the growth of these tumors in the presence of tamoxifen. We conclude that tamoxifen-refractory tumors are not cross-resistant to GW5638, despite the structural similarities of the two compounds. These data support the therapeutic potential of GW5638 for the treatment of tamoxifen-resistant breast cancers.</p>					
<b>14. SUBJECT TERMS</b> Breast Cancer estrogen receptor transcription antiestrogens hypothalamic-pituitary axis promoter tamoxifen				<b>15. NUMBER OF PAGES</b> 15	
				<b>16. PRICE CODE</b>	
<b>17. SECURITY CLASSIFICATION OF REPORT</b> Unclassified	<b>18. SECURITY CLASSIFICATION OF THIS PAGE</b> Unclassified	<b>19. SECURITY CLASSIFICATION OF ABSTRACT</b> Unclassified	<b>20. LIMITATION OF ABSTRACT</b> Limited		

## FOREWORD

Opinions, interpretations, conclusions and recommendations are those of the author and are not necessarily endorsed by the U.S. Army.

N/A Where copyrighted material is quoted, permission has been obtained to use such material.

N/A Where material from documents designated for limited distribution is quoted, permission has been obtained to use the material.

see Citations of commercial organizations and trade names in this report do not constitute an official Department of Army endorsement or approval of the products or services of these organizations.

N/A In conducting research using animals, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and use of Laboratory Animals of the Institute of Laboratory Resources, national Research Council (NIH Publication No. 86-23, Revised 1985).

✓ For the protection of human subjects, the investigator(s) adhered to policies of applicable Federal Law 45 CFR 46.

N/A In conducting research utilizing recombinant DNA technology, the investigator(s) adhered to current guidelines promulgated by the National Institutes of Health.

N/A In the conduct of research utilizing recombinant DNA, the investigator(s) adhered to the NIH Guidelines for Research Involving Recombinant DNA Molecules.

N/A In the conduct of research involving hazardous organisms, the investigator(s) adhered to the CDC-NIH Guide for Biosafety in Microbiological and Biomedical Laboratories.

Caroline E. Connor 6/23/99  
PI - Signature Date

## TABLE OF CONTENTS

I.	Front Cover	Page 1
II.	Report Documentation Page	Page 2
III.	Foreword	Page 3
IV.	Table of Contents	Page 4
V.	Introduction	Pages 5
VI.	Body	Pages 6-10
VII.	References	Page 11
VIII.	Appendix	Pages 12-15

## INTRODUCTION


Estrogen mediates its biological responses through a specific interaction with the estrogen receptor (ER), an intracellular transcription factor which can bind specific DNA sequences and regulate gene expression (1). SERMs (selective estrogen receptor modulators) also bind to ER but induce unique conformations in the protein, allowing for tissue-selective effects (2, 3). One such compound, tamoxifen, has proven clinically useful in the treatment of ER-positive breast cancer. However, despite its bone-protective capabilities, tamoxifen has undesirable uterotrophic effects and, during treatment of breast cancer, resistance to the compound eventually develops (4). Therefore, there is a clinical need for improved SERMs in breast cancer therapy and prevention. GW5638, a compound which has been shown to be bone-protective and non-uterotrophic in animal models, appears to be mechanistically distinct from tamoxifen in a variety of *in vitro* and *in vivo* assays (5). This suggests that GW5638 may offer utility against tamoxifen-refractory breast cancers. The goal of this project was to determine if GW5638 could inhibit the growth of tamoxifen-resistant tumors in an athymic mouse model. Additionally, in an effort to understand the property common to estrogens and SERMs that permits their action as bone-protective agents, we sought to identify GW5638-responsive genes in bone.

## BODY

### A. Tumor Inhibition Studies

The MCF-7 breast cancer model is well-established in athymic mice. The cell line is estrogen-dependent and tumors are growth inhibited by tamoxifen (6, 7). For these experiments, ovariectomized (OVX) female mice are given slow-release estrogen pellets prior to tumor implantation. We first evaluated the potential of GW5638 to inhibit the growth of MCF-7 tumors. Experimental groups were as follows: 1) placebo injections, 2) tamoxifen injections, or 3) GW5638 injections. Similar to tamoxifen, GW5638 inhibited tumor growth (Figure 1). We are currently repeating this experiment to verify the results, confident that GW5638 has inhibitory effects.


Figure 1.



Tamoxifen, the standard adjuvant therapy for estrogen receptor (ER)-positive breast cancer, offers initial success but eventually leads to resistance in many patients (8). This paradigm is also seen in the athymic mouse model, in which other laboratories have shown that continued treatment with tamoxifen leads to a tumor which becomes dependent upon rather than inhibited by the drug (9). One of the goals of this project was to assess whether GW5638 could inhibit the growth of tumors no longer sensitive to tamoxifen inhibition. In order to complete this aim, we first had to develop and characterize a series of tamoxifen-resistant tumors. This was accomplished over the course of several years through continuous dosing with tamoxifen and repeated tumor passaging from animal to animal.

Once tumors began to consistently grow in the presence of tamoxifen, they developed an estrogen-independent phenotype, unlike the parental MCF-7 tumor. This phenomenon is consistent with previous results from other laboratories (9). When analyzed in athymic mice supplied by two different companies, animals with putative tamoxifen-resistant tumors were treated with and without tamoxifen for 5 weeks, as seen in Figure 2. Tumors grew only in animals dosed with tamoxifen; we had created the expected tamoxifen-dependent tumor variant.

**Figure 2.**




Once the tamoxifen-dependent tumors were characterized, we tested the ability of GW5638 to inhibit their growth. Initiation of experiments was difficult due to low tumor take rate and extremely slow growth rate of these tumors as compared to original MCF-7 tumors. Animals were treated with tamoxifen injections three times weekly to establish tumor growth, at which time animals were randomized by tumor volume into three groups. The first treatment group was given injections of both tamoxifen and oil to demonstrate the growth of tumors in the presence of tamoxifen. The second group was given injections of GW5638 and oil to determine whether the tumors could be stimulated with GW5638. We predicted that GW5638 would not stimulate tumor growth, as we believe this compound is mechanistically distinct from tamoxifen. The third group was given injections of both tamoxifen and GW5638 to test for the ability of GW5638 to inhibit the growth of tamoxifen-stimulated tumors. This experiment was performed initially in OVX athymic mice and the results demonstrate that a) GW5638 does not stimulate the growth of the



tamoxifen-dependent tumors and b) GW5638 can inhibit the growth of such tumors in the presence of tamoxifen (Figure 3).

Figure 3.


### Inhibition of Tamoxifen-resistant Tumors by GW5638 in OVX mice



To verify this experiment, we repeated it in non-OVX mice, a more clinically relevant model, and obtained similar results, as shown in Figure 4.

Figure 4.


### Inhibition of Tamoxifen-resistant Tumors in Non-OVX Mice



During the months of development of the tamoxifen-resistant tumors, we tested the effect of GW5638 on the MCF-7/LCC2 cell line *in vivo*. This cell line, derived *in vitro*

from the parental MCF-7 cell line at the Lombardi Cancer Center, is reported to be hormone-independent, sensitive to estrogen, and resistant to the growth inhibitory effects of tamoxifen when analyzed in the athymic mouse model (8). We performed a tumor inhibition experiment comparing treatments with placebo, estrogen, tamoxifen, and GW5638. As demonstrated in Figure 5, in our experiment, the tumor variant grew equally well in the absence or presence of estrogen and both SERMs tested were able to inhibit its growth. This suggests that GW5638 can inhibit the growth of an estrogen-independent tumor.

Figure 5.



#### B. Identification of GW5638-Responsive Genes

Despite the observation that all ER ligands, except pure antiestrogens, are protective against bone loss, specific genes which mediate this effect have not been identified (10). We attempted to identify GW5638-regulated genes in bone using differential display PCR (ddPCR) (11, 12). Female rats were OVX at 6 months of age, and animals were treated with either estrogen, tamoxifen, GW5638, or vehicle on day 15 post-OVX. Total RNA from pulverized bone was pooled for each treatment group. This RNA was reverse transcribed into partial cDNA sequences through PCR amplification, prior to being used in extensive ddPCR analysis. Bands which appeared to be uniquely regulated in several repeated

expression may have been hard to detect when using RNA blots composed of a variety of different bone cell types.

Due to our interest also in identifying genes regulated by estrogen itself in bone, we sent RNA samples from vehicle- or estrogen-treated animals to Genome Systems, Inc. An analysis was performed using this company's verified human microarray, consisting of approximately 7,000 clones, in hopes of finding regulated genes. Only two genes were shown to be regulated by estrogen treatment; however, the subsequent Northern analysis could not detect any changes in RNA levels.

Our difficulty in identifying estrogen- or GW5638-regulated genes in bone was disappointing. However, one possible explanation is that GW5638 may not, in fact, regulate a unique subset of genes. Additionally, the fact that many classes of SERMs are bone-protective may be due to non-genomic effects of these compounds, which would not have been seen in the aforementioned analyses.

## **Conclusions**

We conclude that GW5638 is able to inhibit the growth of MCF-7 breast cancer tumors in mice. Additionally, we have created a tamoxifen-resistant tumor variant which is dependent upon tamoxifen for growth. GW5638, a mechanistically distinct compound, can inhibit the growth of such tumors and does not stimulate these tumors when given alone. Unfortunately, our efforts to identify GW5638-responsive genes were ineffective; this is not surprising as specific SERM-regulated genes have not yet been identified, perhaps indicating that non-genomic effects may play a role in mediating bone-protection.

## REFERENCES

1. Evans, R. M. The Steroid and Thyroid Hormone Receptor Superfamily, *Science*. 240: 889-895, 1988.
2. McDonnell, D. P., Clemm, D. L., Hermann, T., Goldman, M. E., and Pike, J. W. Analysis of Estrogen Receptor Function *in Vitro* Reveals Three Distinct Classes of Antiestrogens, *Mol. Endocrinol.* 9: 659-669, 1995.
3. Smith, C. L., Nawaz, Z., and O'Malley, B. W. Coactivator and Corepressor Regulation of the Agonist/Antagonist Activity of the Mixed Antiestrogen, 4-Hydroxytamoxifen, *Mol. Endocrinol.* 11: 657-666, 1997.
4. Kedar, R. P., Bourne, T. H., Powles, T. J., Collins, W. P., Ashley, S. E., Cosgrove, D. O., and Campbell, S. Effects of tamoxifen on uterus and ovaries of postmenopausal women in a randomised breast cancer prevention trial, *Lancet*. 343: 1318-1321, 1994.
5. Willson, T. M., Norris, J. D., Wagner, B. L., Asplin, I., Baer, P., Brown, H. R., Jones, S. A., Henke, B., Sauls, H., Wolfe, S., Morris, D. C., and McDonnell, D. P. Dissection of the Molecular Mechanism of Action of GW5638, a Novel Estrogen Receptor Ligand, Provides Insight into the Role of Estrogen Receptor in Bone, *Endocrinology*. 138: 3901-3911, 1997.
6. Giovenella, B. P., Stehlin, J. S., Williams, J., L.J., Lee, S.-S., and Shepard, R. C. Heterotransplantation of Human Cancers into Nude Mice, *Cancer*. 42: 2269-2281, 1978.
7. Soule, H. D. and McGrath, C. M. Estrogen Responsive Proliferation of Clonal Human Breast Carcinoma Cells in Athymic Mice, *Cancer Lett.* 10: 177-189, 1980.
8. Brunner, N., Frandsen, T. L., Holst-Hansen, C., Bei, M., Thompson, E. W., Wakeling, A. E., Lippman, M. E., and Clarke, R. MCF7/LCC2: A 4-Hydroxytamoxifen Resistant Human Breast Cancer Variant That Retains Sensitivity to the Steroidal Antiestrogen ICI 182,780, *Cancer Res.* 53: 3229-3232, 1993.
9. Gottardis, M. M. and Jordan, V. C. Development of Tamoxifen-stimulated Growth of MCF-7 Tumors in Athymic Mice after Long-Term Antiestrogen Administration, *Cancer Res.* 48: 5183-5187, 1988.
10. McDonnell, D. P. and Norris, J. D. Analysis of the Molecular Pharmacology of Estrogen Receptor Agonists and Antagonists Provides Insights into the Mechanism of Action of Estrogen in Bone, *Osteoporosis Int.*, 1997.
11. Liang, P. and Pardee, A. B. Differential Display of Eukaryotic Messenger RNA by Means of the Polymerase Chain Reaction, *Science*. 257: 967-971, 1992.
12. Liang, P., Averboukh, L., and Pardee, A. B. Distribution and cloning of eukaryotic mRNAs by means of differential display: refinements and optimization, *Nucleic Acids Res.* 21: 3269-3275, 1993.

## APPENDIX

### *1) Key Research Accomplishments*

- Demonstrated the ability of GW5638 to inhibit the growth of MCF-7 breast cancer tumors in athymic mice.
- Developed tamoxifen-resistant tumors *in vivo*.
- Demonstrated the ability of GW5638 to inhibit the growth of tamoxifen-resistant breast cancer tumors in athymic mice.

### *2) Reportable Outcomes*

#### Manuscripts/Abstracts/Presentations

1. **Connor, C.E.**, Norris, J.D., Gottardis, M.M., Willson, T.M., Dewhirst, M.W., and McDonnell, D.P. Inhibition of tamoxifen-refractory breast cancer by GW5638. (1999) *In preparation*.
2. **Connor, C.E.**, Norris, J.D., Dewhirst, M.W., and McDonnell, D.P. Therapeutic Potential of GW5638 for the Treatment of Tamoxifen-Resistant Breast Cancer. (1999) SPORE Investigators Workshop, Rockville, MD *Selected talk*.
3. **Connor, C.E.**, Buxton, J., Dewhirst, M.W., and McDonnell, D.P. Evaluation of GW5638, a novel antiestrogen, as an antitumor agent in nude mice. (1997) 20th Annual San Antonio Breast Cancer Symposium, San Antonio, TX *Poster presentation*.
4. **Connor, C.E.**, Buxton, J., Dewhirst, M.W., and McDonnell, D.P. Evaluation of GW5638, a novel antiestrogen, as an antitumor agent in nude mice. (1997) SPORE Investigators Workshop, Rockville, MD *Poster presentation*.

#### Patents

1. McDonnell, D.P., Norris, J.D., **Connor, C.**, and Wijayaratne, A. (1998) A method of preventing or treating estrogen-dependent diseases and disorders. (pct. app.)

#### Development of tumor line

1. Development of tamoxifen-dependent MCF-7 tumor variants in athymic mice.

### *3) Cited Abstracts (attached)*

## **SPORE Investigators Workshop, 1999**

### **Therapeutic Potential of GW5638 for the Treatment of Tamoxifen-Resistant Breast Cancer**

Caroline E. Connor,<sup>1</sup> John D. Norris, Ph.D.,<sup>1</sup> Mark W. Dewhirst, Ph.D., D.V.M.,<sup>2</sup> and Donald P. McDonnell, Ph.D.<sup>1</sup>

<sup>1</sup>Department of Pharmacology and Cancer Biology, <sup>2</sup>Department of Radiation Oncology  
Duke University Medical Center, Durham, NC 27710

SERMs (selective estrogen receptor modulators) are believed to regulate estrogen receptor (ER) transcriptional activity by inducing distinct conformational changes in the ligand-binding domain of the receptor, each of which has a unique biological consequence. Tamoxifen is the most widely used SERM in the treatment of breast cancer; however, its therapeutic utility is limited by the eventual development of resistance. We believe that this may result from the ability of tamoxifen to switch from an antagonist to a partial agonist in breast cancer cells. We hypothesize that compounds which are mechanistically distinct from tamoxifen may be able to inhibit the growth of tamoxifen-resistant tumors. Toward this goal, our laboratory has identified GW5638, a SERM which lacks partial agonist activity. Additionally, using phage display technology, we identified peptides which bind to ER specifically in the presence of tamoxifen and not GW5638. This demonstrates that the two compounds expose unique surfaces on ER, attesting to their mechanistic differences. These data provide a rationale for the examination of GW5638 as a treatment for tamoxifen-refractory breast cancer tumors.

For our pre-clinical evaluation of GW5638, we developed several MCF-7 cell breast cancer tumor variants in mice which are resistant to the tumorigenic effects of tamoxifen. Unlike the parent tumor line, these tumors actually require tamoxifen, but not estrogen, for growth. The resistant tumors were implanted into ovariectomized (OVX), athymic mice and animals were treated with tamoxifen alone, GW5638 alone, and a combination of tamoxifen and GW5638 at equal doses. Two important findings emerged from these studies: 1) GW5638 alone cannot support the growth of tamoxifen-dependent tumors, and 2) GW5638 inhibits the growth of these tumors in the presence of tamoxifen. These results were confirmed in a similar experiment conducted in non-OVX mice, a more clinically relevant paradigm. We conclude that tamoxifen-refractory tumors are not resistant to GW5638, despite the structural similarities of the two compounds. Cumulatively, these data support the introduction of GW5638 into the clinic for the treatment of tamoxifen-resistant breast cancers.

## 20th Annual San Antonio Breast Cancer Symposium, 1997

EVALUATION OF GW5638, A NOVEL ANTIESTROGEN, AS AN ANTITUMOR AGENT IN NUDE MICE. Connor CE\*, Buxton J, Dewhirst, MW, and McDonnell DP, Duke University Medical Center, Durham, North Carolina 27710.

Using the athymic nude mouse model, it has been previously demonstrated that tamoxifen is able to inhibit the growth of MCF-7 breast cancer tumors. Continued treatment with the drug resulted in resistance which actually manifests as tamoxifen dependence. GW5638 is a recently developed tamoxifen analog which offers bone and cardiovascular protection. Interestingly, this compound not only fails to demonstrate uterotrophic activity in ovariectomized rats but also opposes the partial agonist activity of tamoxifen in all cell and promoter contexts examined, suggesting that it is mechanistically distinct from previously studied antiestrogens. These unique characteristics make GW5638 a likely candidate for the treatment of tamoxifen-failed breast cancer patients. *In vivo* experiments were conducted to determine whether this compound would behave as tamoxifen in the nude mouse breast cancer model. A dose-response experiment revealed that at equivalent doses, GW5638 suppressed tumor growth as well as tamoxifen. To determine whether this compound can rescue tamoxifen-failed tumors, efforts are underway to provoke failure through long-term tamoxifen treatment. An alternative approach has been to use MCF-7/LCC2, a cell line variant which no longer responds to tamoxifen treatment. Previous studies have indicated that only ICI, the pure steroidal antagonist, can inhibit the growth of this cell line *in vitro*. Our current studies will allow us to ascertain if GW5638 can inhibit growth of the tumor *in vivo*.

We conclude that GW5638 is as efficacious as tamoxifen in its ability to inhibit the growth of MCF-7 breast cancer tumors in nude mice and may be able to rescue tamoxifen-failed tumors.

## **SPORE Investigators Workshop, 1997**

### **EVALUATION OF GW5638, A NOVEL ANTIESTROGEN, AS AN ANTITUMOR AGENT IN NUDE MICE**

Caroline E. Connor, Jake Buxton, Mark W. Dewhirst, and Donald P. McDonnell  
Duke University Medical Center, Durham, NC

Estrogen mediates its biological responses through a specific interaction with the estrogen receptor (ER), an intracellular protein located within target cell nuclei. Following binding to latent receptor, hormone promotes conversion to a transcriptionally active form which can bind specific DNA sequences and regulate gene expression. Antiestrogens, despite their ability to deliver ER to DNA, freeze the receptor in different conformational states, all of which demonstrate reduced transcriptional activity in classical ER response systems. Recent evaluations of the biological activity of ER antagonists *in vitro* have led to the classification of antiestrogens into three distinct groups represented by tamoxifen, keoxifene, and ICI. Thus, a link was established between receptor structure and biological activity.

The partial agonist 4-OH tamoxifen is widely used as adjuvant therapy for the treatment of breast cancer. Tamoxifen enigmatically exhibits desirable antagonist activity in breast, partial agonist activity in the cardiovascular system and bone, but is unfortunately associated with uterine hypertrophy. GW5638 is a recently developed tamoxifen analog which offers bone and cardiovascular protection. Interestingly, this compound not only fails to demonstrate uterotrophic activity in ovariectomized rats but also opposes the partial agonist activity of tamoxifen in all cell and promoter contexts examined, suggesting that it is mechanistically distinct from previously studied antiestrogens. These unique characteristics make GW5638 a likely candidate for the treatment of tamoxifen-failed breast cancer patients.

Using the athymic nude mouse as a model, we have demonstrated that GW5638 is able to potently inhibit the growth of MCF-7 breast cancer tumors. To determine whether this compound can rescue tamoxifen-failed tumors, efforts are underway to provoke failure through long-term tamoxifen treatment. An alternative approach has been to use MCF-7/LCC2, a cell line variant which no longer responds to tamoxifen treatment. Previous studies have indicated that only ICI, the pure steroidal antagonist, can inhibit the growth of this cell line *in vitro*. Our current studies will allow us to ascertain if GW5638 can inhibit growth of the tumor *in vivo*. Future work involves the identification of genes regulated by this compound as a means of understanding its tissue-selective action.





DEPARTMENT OF THE ARMY  
US ARMY MEDICAL RESEARCH AND MATERIEL COMMAND  
504 SCOTT STREET  
FORT DETRICK, MARYLAND 21702-5012

*Rec'd*  
*7/19/2000*

REPLY TO  
ATTENTION OF:

MCMR-RMI-S (70-1y)

6 Jul 00

MEMORANDUM FOR Administrator, Defense Technical Information  
Center, ATTN: DTIC-OCA, 8725 John J. Kingman  
Road, Fort Belvoir, VA 22060-6218

SUBJECT: Request Change in Distribution Statements

1. The U.S. Army Medical Research and Materiel Command has  
reexamined the need for the limitation assigned to technical  
reports written for the following awards:

DAMD17-94-C-4068	ADB218322
DAMD17-94-V-4036	ADB232944
DAMD17-94-J-4481	ADB222571
DAMD17-95-C-5054	ADB227112
DAMD17-96-1-6016	ADB228823
DAMD17-96-1-6073	ADB248567
DAMD17-94-J-4057	ADB221437, ADB247857
DAMD17-96-1-6069	ADB230256, ADB239319
DAMD17-95-1-5067	ADB236775, ADB249592
DAMD17-94-J-4308	ADB225776, ADB234457, ADB249935
DAMD17-96-1-6087	ADB232086, ADB238945, ADB250354
DAMD17-96-1-6075	ADB228777, ADB238338, ADB249653
DAMD17-95-1-5008	ADB225250, ADB236089, ADB243691
DAMD17-94-J-4310	ADB222453, ADB235860, ADB247801

Request the limited distribution statement for Accession Document  
Numbers be changed to "Approved for public release; distribution  
unlimited." These reports should be released to the National  
Technical Information Service.

2. Point of contact for this request is Ms. Virginia Miller at  
DSN 343-7327 or by email at Virginia.Miller@det.amedd.army.mil.

FOR THE COMMANDER:

*Phyllis Rinehart*  
PHYLLIS M. RINEHART  
Deputy Chief of Staff for  
Information Management